



**ECOSUR**

# **El Colegio de la Frontera Sur**

**Descomposición de hojarasca de especies arbóreas en tres  
fases sucesionales de la Selva Mediana, Campeche, México**

**TESIS**

**Presentada como requisito parcial para optar al grado de Maestría en Ciencias**

**por**

**Francisco Javier Xuluc Tolosa**

**2001**

877  
BCLC  
DONACION  
04-09-01



**ECOSUR**

## El Colegio de la Frontera Sur

Descomposición de hojarasca de especies arbóreas en tres  
fases sucesionales de la Selva Mediana, Campeche, México

**TESIS**

Presentada como requisito parcial para optar al grado de Maestría en Ciencias

por

**Francisco Javier Xuluc Tolosa**

TE/631.4/X8/

Xuluc Tolosa, Francisco Javier

Descomposición de hojarasca de especies

TE 540

**18 DIC. 2001**

## AGRADECIMIENTOS

Al Colegio de la Frontera Sur (ECOSUR) por el otorgamiento de la beca para estudios de posgrado con recursos de la SEP/CONACYT.

Al Consejo Tutelar integrado por el Dr. Henricus Franciscus M. Vester; por los M. en C. Neptalí Ramírez Marcial y Jorge Castellanos Albores.

A Deborah Lawrence por su asesoría y revisión del manuscrito.

A Humberto Bahena Basave por su trabajo fotográfico empleado en la presentación y su apoyo incondicional en el campo.

A Margarito Tuz Novelo por su apoyo incondicional en la colecta de hojarasca.

A Diego Pérez Salicrup por los datos proporcionados de las parcelas.

Sophie Calme y Mauro Sanvicente por su hospitalidad en Zoh Laguna.

Jesús Carmona y Juan Morales López por la realización de los análisis químicos de la hojarasca.

Noemí Salas por su ayuda en la identificación de la fauna registrada en la hojarasca.

Patricia Montañez por su intercambio de experiencias respecto al tema tratado.

A mi familia que me ha apoyado siempre e incondicionalmente y en especial a mis padres Javier Xuluc Manzanilla y Berta Tolosa Canché por su amor y confianza.

A mis compañeros por haberme permitido intercambiar experiencias durante los cursos

## CONTENT

Abstract.....	1
Introduction .....	2
Study area.....	3
Methods .....	5
Forest stand selection.....	5
Leaf litter decomposition of tree species.....	6
<i>In situ</i> decomposition of local leaf litter .....	7
Nutrient concentration.....	8
Statistical analysis .....	8
Results .....	10
Leaf litter decomposition of tree species.....	10
<i>In situ</i> decomposition of local leaf litter .....	12
Discussion.....	13
Leaf litter decomposition of individual species.....	13
<i>In situ</i> decomposition of local leaf litter .....	17
Conclusion .....	18
References.....	19

## ABSTRACT

Leaf litter decomposition rate and nutrient concentration (N, C, P) of three species: *Croton lundellii* Standl., *Metopium brownei* (Jacq.) Urban and *Manilkara zapota* (L.) van Royen) were evaluated in different phases of development (3, 13 and > 50 years old) of secondary dry forest in Campeche, Mexico, to test two hypotheses: 1) leaf decomposition rate is higher in older successional stages of secondary dry forest, and 2) decomposition rate is more rapid in pioneer species than late successional species, because of differences in substrate quality. The litterbag method was employed to evaluate decomposition. Decomposition was related to forest development phase but the strongest influence was due to leaf chemical composition. Decomposition was fastest in *C. lundellii*, intermediate in *M. brownei* and slowest in *M. zapota*. The C/N, C/P ratios and total N and P concentrations were good indicators of decomposition rate across species in all successional stages. In comparing leaf litter collected locally and decomposed in situ, the earlier successional phase showed higher decomposition rates than later stages of succession. Neither N concentration or C/N ratio explained this difference, but it was correlated with higher P concentration.

**Keywords:** *Croton lundellii*, *Metopium brownei*, *Manilkara zapota*, litter quality, decay rate, succession.

## INTRODUCTION

The leaf litter deposition and decomposition are recognized as critical pathways of organic matter and nutrient flux in tropical dry forest systems (Melillo *et al.*, 1982; Proctor, 1983; Alvarez *et al.*, 1992). The leaf litter aboveground is the main input of nutrients to the soil (Herborn and Congdon, 1998; Martinez-Yrizar, 1995; Montañez, 1998). The production and decomposition of leaf litter in dry forests are influenced by the seasonality of precipitation and plant composition (Anderson and Swift, 1983; Cuevas, 1995, Lawrence, submitted). During the wet season, conditions for growth and activity of soil microorganisms are improved, thus increasing decomposition rate. Studies on soil invertebrates show that during the dry season the soil fauna moves deeper in the soil (Prieto *et al.*, 1999). Due to this difference between dry and wet season, in some tropical dry forests an inhibition of decomposer community results in a transient accumulation of litter on the forest floor during the dry season (Swift and Anderson, 1989).

In dry forests on the Yucatan Peninsula, litter production (Whigham *et al.*, 1990; Whigham *et al.*, 1991) and decomposition of fine and coarse woody detritus (Harmon *et al.*, 1995) have been studied, but leaf litter decomposition along a successional gradient was not investigated. Montañez (1998) found that leaf litter decomposition of tree species in home gardens depended on season and species, where a slow decomposition occurred during the dry season and fast during the rainy season. Different decomposition rates among tropical species have been documented (Babbar and Ewel, 1982; Alvarez and Becerra, 1996; Montañez,

1998; Sundarapandian & Swamy, 1999) and leaf quality, especially the C/N ratio, has been proposed as a sound predictor of decomposition rate (Heal *et al.*, 1997). Because the floristic composition and forest structure changes during secondary succession (Brown and Lugo, 1990), decomposition will depend on the age of the forest (Coleman and Crossley, 1996). If pioneer species have high decomposition rates due to higher leaf nutrient content, then slower decomposition can be expected in mature forests. On the other hand, because in old-growth forests the ecosystem as an integrated system may be more efficient in the recycling of nutrients (Brown and Lugo, 1990), we may expect higher decomposition rates in older forests.

The major problem in dry forest is land use conversion, which entails disruption of ecological processes like productivity, hidrological process, soil erosion and nutrient cycling, and biodiversity loss (Mass, 1995).

The objective of this study was to evaluate the leaf litter decomposition rate of tree species in three different growth phases of tropical dry secondary forest (3, 13 and > 50 years old). Two hypotheses were tested: first, that decomposition rate is higher in older successional stages of secondary dry forest, and second, that it is higher in pioneer species relative to late successional species due to differences in substrate quality.

## **STUDY AREA**

The study was carried out on the rim of the Calakmul biosphere reserve in the state of Campeche, southern Mexico. The reserve is situated between 17° 45' and



19° 15' N, 89° 15' and 90° 10' W. This zone comprises one of the largest well-preserved tropical dry forest in southern Mexico. Approximately 34% of the total area (723,185 ha) corresponds to the core area, which is surrounded by a human populated buffer zone (Díaz, 2000). About 50,000 inhabitants coming from 26 states of the Mexican Republic have been settled in 120 villages since 1970's. The community of "El Refugio" (northeastern part of the Reserve, 107 inhabitants) was chosen because there are a number of permanent plots established by the Land cover and Land use Change project of Clark University, Harvard forest and ECOSUR financed by NASA (Turner *et al.*, 2001).

Topography is relatively flat and characterized by a karstic plateau, reaching a maximum elevation of 380 m above sea level. This plateau shows an altitudinal gradient, decreasing from South to North in the reserve (García and March, 1990). The soils are classified as rendzina, gleysol, litosol and vertisol with different kinds of texture (FAO, 1970; García and March, 1990). The climate according to the Köppen system modified for García (1973) is seasonally dry subhumid (Aw) with a summer rainy season (June-October). The annual precipitation is around 945 mm, and the mean annual temperature is ca. 22° C. The dry season extends between February and May. In August, there is a short dry season (locally called "canicula"). Monthly rainfall and temperature distribution for the study period are presented in Fig. 1.

The vegetation types in the area correspond to tropical semi-evergreen forest (*selva mediana subperennifolia*), tropical low semi-evergreen forest (*selva baja subperennifolia*) and secondary forest (Lundell, 1934; Miranda, 1958, Rzedowski, 1981; Flores *et al.*, 1990, Flores and Espejel, 1994). These types are classified as

tropical dry forest in the Holdridge system (Murphy and Lugo, 1986). The forest in this study is characterized by the presence of both evergreen and deciduous species. Canopy height is between 15 and 25 m. The characteristic tree species are *Manilkara zapota* (L.) van Royen, *Vitex gaumeri* Greenman, *Lysiloma latisiliquum* (L.) Benth., *Bursera simaruba* (L.) Sarg., *Metopium brownei* (Jacq.) Urban, *Brosimum alicastrum* Swartz, *Piscidia piscipula* (L.) Sarg., *Malmea depressa* (Baill) R. E. Fries, *Guetarda combsii* Urban, *Thouinia paucidentata* Radlk., *Thevetia gaumeri* Hemsley, *Lonchocarpus rugosus* Benth., *Croton lundellii* Standl., and *Cecropia obtusifolia* Berthol.

## **METHODS**

### **Forest stand selection**

During 1999, a number of permanent circular plots (500 m<sup>2</sup>) were established in secondary forest stands of known age, all of which have been used for slash and burn agriculture. Pérez (pers. com.) described the floristic composition and forest structure in these permanent plots and Lawrence (submitted) determined the litterfall dynamics and soil physical and chemical characteristics. Three successional stages with different environmental conditions were chosen: (1) secondary forests 3 years old reaching 2-3.5 m in height dominated by young trees, shrubs (> 1 cm and < 4.5 cm dbh) and climber plants; (2) secondary forests 13 years old with a high density of trees (> 4.5 cm and < 16 cm dbh) and shrubs 5-8 m tall; and (3) mature forest more than 50 years old, dominated by tall trees (>30 cm dbh) reaching 13-16 m in height (Pérez, pers. com.). Two plots of each

successional phase were chosen from the plots formerly established in the litterfall collection experiment.

### **Leaf litter decomposition of tree species**

A leaf litter decomposition experiment was carried out during a period of seven months from April to October, 2000. Three plant species were used in the experiment: *Croton lundellii* Standl., an early successional deciduous species; *Metopium brownei* (Jacq.) Urban; a late successional deciduous species, and *Manilkara zapota* (L.) van Royen, an evergreen late successional species. These species were typical components of forest composition and structure in the study area (Perez, pers. com.). Entire leaves of *C. lundellii* and *M. brownei* were harvested directly from adult trees during February and March, just before abscission. Then they were oven dried at 60 °C for 72 hours. Recently fallen leaves of *M. zapota* were collected under the trees, taking care that all leaves were in a similar condition. Leaf litter was used to determine decomposition because this plant organ represents a substantial portion (50 to 80%) of the total aboveground litter production in terms of biomass (Toky and Ramakrishnan, 1983a; Alvarez *et al.*, 1992; Martinez-Yrizar, 1995; Montañez, 1998; Sundarapandian and Swamy, 1999). Because of their relatively higher nutrient concentration, they represent the major nutrient input as well nutrients (Toky and Ramakrishnan, 1983b; Alvarez *et al.*, 1992). Furthermore, leaves decompose faster than twigs and other woody materials (Anderson and Swift, 1983), thus providing rapid inputs to the soil nutrient cycle.

Leaf decomposition was evaluated through the litterbag technique described by Swift *et al.* (1979) and Alvarez *et al.* (1992). This technique has proven useful in delineating and analyzing differences in decomposition rate (Coleman and Crossley, 1996). Mesh bags 15 x 20 cm in size with 2 mm mesh were filled with 30 g of dried leaves from one of the species (Montañez, 1998). The leaves of an individual species were mixed to obtain a uniform mixture before filling the mesh bags. The 1-2 mm mesh size was considered sufficiently small to prevent losses of litter due to breakage, but sufficiently large to permit the access of decomposers (Sundarapandian and Swamy, 1999). On April 1, three replicates of six litterbags per species were labeled and placed on the forest floor at random on the border of each plot. A total of 54 leaf litterbags were set out in each plot (108 in each successional phase), with 324 bags for the entire study.

### ***In situ* decomposition of local leaf litter**

In order to compare the decomposition of the total litter deposition and decomposition among different successional stages, bags containing local leaf litter were deployed as well. These bags were filled with 30 g of the leaf fraction, obtained from litterfall harvested in four traps (1x1 m<sup>2</sup>) during February and March. An additional 18 mesh bags of local leaf litter were placed in each plot (36 per successional stage). A total of 108 bags were employed for the study of *in situ* decomposition of local leaf litter.

At six time intervals (35, 63, 91, 125, 156 and 190 days after start of experiment) three bags per species and three of the local leaf litter samples were retrieved from the plots for analysis. The analysis consisted of measuring the mass of leaves left

in the bag after drying in paper bags in an oven at 60 °C for 72 hours. Differences between the mass at the beginning of the experiment and that at the different time periods were calculated to evaluate leaf decomposition (Singh and Gupta, 1977).

### **Nutrient concentration**

Chemical analyses were undertaken to determine leaf litter quality for each species and local leaf litter sample, and to track changes in leaf litter quality during the experiment. Total C, N, and P concentration were determined in subsamples from each litterbag before the experiment, and each time a bag was taken out for measurement of remaining mass. After dry weight was measured, the leaves in all three bags per species were mixed and milled to generate homogeneous, composite samples for the analysis. Determination of N and P was done using a modified Kjeldahl method (Harris, 1970) followed by photometric analysis (AOAC, 1994). Total organic matter was estimated using the Walkley and Black method (Anderson and Ingram, 1993). Organic carbon contents were calculated from the organic matter values using a conversion factor of 0.58 g carbon per gram organic matter.

### **Statistical analysis**

To describe and compare the decomposition rates between ages of secondary forest development and among tree species, a repeated-measures analysis of variance (Ende, 1993;  $p < 0.10$ ) was used with fixed factors and split-plot nested design. The plots were the subjects; time was the within-subject factor, and

treatment (tree species nested in successional stages) the between-subject factor. This analysis took into account the correlation between dates and allowed comparison and fitting of the response patterns (shape of the curve). We obtained the Huynh-Feldt corrected probability (Ende, 1993) to decide rejection or acceptance of the differences with time and the time by treatment interactions ( $p < 0.10$ ). A similar analysis was employed to compare leaf quality between species nested in environmental condition (successional stages). The same procedure was applied to compare the local leaf litter loss and leaf quality among stages of secondary forest development, followed by a Tukey test (Steel and Torrie, 1988). A linear model was applied to calculate annual decay rate constants ( $k$ ) from dry mass remaining for each tree species and for the local leaf litter sample (Wieder and Lang, 1982).

The equation was  $X_t = -k \cdot t + X_0$

where  $X_t$  = remaining mass (g) at time  $t$  (days),  $X_0$  = initial leaf mass, and  $k$  = decomposition rate constant ( $\text{yr}^{-1}$ ). The linear model is defined such that the absolute decomposition rate is constant throughout decomposition, while the relative decomposition rate increases with time (Wieder and Lang, 1982). This model can be applied as well when leaf litter decomposition decreases slowly (Montañez, 1998). The correlation between litter quality and annual decay rate constant was realized based on mean values of C/N ratio, C/P ratio, N and P concentration obtained during the six measurements.

## RESULTS

### Leaf litter decomposition of tree species

The analysis showed no significant ( $p > 0.1$ ) overall effect of successional stage on decomposition (Table 1) but species was a strong determinant of decomposition rate ( $F_{2, 3} = 132.7$ ;  $p = 0.001$ ) (Fig. 2). There is evidence that the decomposition was not constant in time ( $F_{5, 15} = 277.5$ ;  $p < 0.001$ ) and each species varied along time ( $F_{10, 15} = 7.40$ ;  $p < 0.001$ ). Furthermore, decomposition was significantly slightly higher (0.78 g,  $p < 0.010$ ) in the 3 year old forest than in the 13 year old forest.

Leaf litter mass loss for all species followed a similar decomposition pattern through time (Fig. 3a). Decomposition increased with the onset of the rainy season from June to beginning of August (63 to 125 days) then stayed low and relatively constant for the latter part of the study (125 to 190 days). However, *Croton lundellii* started losing mass first, followed by *Metopium brownei* and later by *Manilkara zapota*. *Croton lundellii* showed the greatest mass loss at 190 days (44.3% or 13.3 g of the original mass), *Metopium brownei* showed intermediate loss (30% or 9.0 g), and *Manilkara zapota* showed the lowest mass loss (26.3% or 7.9 g, Fig. 3a).

The C/N ratio followed a similar pattern in all species, declining with time as expected during the process of decomposition (Fig. 3b). By the end of the experiment, *C. lundellii*, which began with a lower C/N ratio, displayed the lowest C/N ratio (mean over all treatments, 36.22). *M. brownei* and *M. zapota*, having lower nutrient quality originally, finished with higher C/N ratios (60.40 and 71.51,

respectively). Although these two species started with a similar C/N ratio, it was much more variable in *M. zapota* than in *M. brownei* throughout the course of the experiment (Fig. 3b). There were significant differences in leaf quality between species ( $F_{2, 3} = 165.4$ ;  $p = 0.000$ ), but not in the way they functioned in different stages of secondary forest development ( $p > 0.1$ ). The species-time interaction was significant, indicating that C/N ratios changed at different rates across species ( $F_{2, 3} = 165.4$ ;  $p = 0.000$ ).

The analysis of nutrients concentration showed a significant effect of species: total N ( $F_{2, 3} = 43.97$ ;  $p = 0.006$ ), total C ( $F_{2, 3} = 86.55$ ;  $p = 0.002$ ) and total P ( $F_{2, 3} = 35.82$ ;  $p < 0.001$ ) but not successional stage ( $p > 0.1$ ). C and N concentrations were significantly different between *C. lundellii* and both *M. zapota* and *M. brownei*; however, *M. zapota* and *M. brownei* were not significantly different. All three species were significantly different in P concentration (Tukey's for all comparisons,  $p < 0.05$ ).

Total nitrogen concentration significantly increased in the leaf tissue of all three species with time ( $F_{6, 12} = 10.68$ ;  $p < 0.001$ ). The highest values were recorded for *C. lundellii* throughout all time intervals. At the end of the experiment, its N concentration (1.35%) was 50% greater than that of *M. brownei* (0.89%) and 90% greater than that of *M. zapota* (0.77%). Conversely, carbon concentration showed no significant trend with time. Through time, *M. zapota* showed higher total C concentration (52.7%), followed by *M. brownei* (51.4%) and *C. lundellii* (45.7%). On the other hand, phosphorus concentration exhibited a notable increase during the first three sampling periods for *C. lundellii* and *M. brownei*, followed by a decline towards the end of the study. *M. zapota* showed a slight increase in the first



months and a subsequent stabilization in P concentration. Phosphorus concentration was highest for *C. lundellii* (0.0381%, averaged over all time intervals), followed by *M. brownei* (0.0359%) and *M. zapota* (0.0345%).

*C. lundellii* had a higher decay rate constant (-2.54) than *M. brownei* (-1.77) and *M. zapota* (-1.46) averaged over all three environmental conditions (all  $R^2$  values > 0.91). Statistical analysis showed that decay rates did not differ significantly ( $p > 0.1$ ) between stages of secondary forest development, although there was a trend toward increasing decomposition rate with forest age (Fig. 2). Differences were recorded between k's species ( $F_{2,3} = 52.36$ ;  $p < 0.005$ ). Tukey test indicated that all species have markedly different decay rates ( $p < 0.05$ ). Annual decay rate was inversely related to C/N and C/P ratios ( $R^2 = -0.829, -0.874$  Fig. 4a, b) and positively correlated with total N and P concentrations ( $R^2 = 0.7486, 0.783$  Fig. 4c,d).

### ***In situ* decomposition of local leaf litter**

Both successional stage ( $F_{2,3} = 11.38$ ;  $p = 0.002$ ) and time ( $F_{2,3} = 75.26$ ;  $p < 0.001$ ) had significant effects on decomposition (Table 2). The mass loss behavior of local leaf litter decomposing *in situ* was similar in all three phases of secondary forest (Fig. 5a). Cumulative mass loss increased through time, and began to level off after 190 days. Mass loss was higher in the early successional phase during all time intervals. Ultimately, leaf litter from 3 year old forests lost 43% (12.9 g), vs. 33% (9.9 g) and 28% (8.4 g) in 13 and > 50 year old forests, respectively.

The C/N ratio diminished significantly ( $F_{6, 12} = 8.79$ ;  $p < 0.001$ ) with time for all successional phases, but did not differ significantly among phases ( $p > 0.1$ , Fig. 5b). C and N concentration did not differ significantly ( $p > 0.1$ ) among stages. Successional stage was a significant factor only in explaining changes in P concentration ( $F_{2, 3} = 18.19$ ;  $p = 0.021$ ). P concentration was moderately higher in early successional forest (0.0382%) than in > 50 year old secondary forest 0.0372% or 13 year old secondary forest (0.0366%). Nutrient dynamics varied depending on the nutrient considered. N concentration increased through time, and C concentration remained steady, whereas P concentration increased initially and then declined.

Estimated annual decay rates were slightly higher ( $k = -2.40$ ) for early than for later successional phases of secondary vegetation (-1.95 and -1.94 for 13 and > 50 year old stands, respectively). Decomposition rates did not vary significantly ( $p > 0.1$ ) between phases of secondary forest development.

## DISCUSSION

### Leaf litter decomposition of individual species

In this study, the results indicate that successional stage does not have a strong effect on decomposition, but the decay rate of individual species associated with those stages varied considerably. The tree species were chosen as representatives of different successional groups: early successional *Croton lundellii*, mid successional *Metopium brownei* and late successional *Manilkara zapota*. Along this sequence, decomposition rate decreased. This pattern can be

explained by indices of leaf litter quality, such as initial N concentration and C/N ratio, which have been widely employed and are relatively easy to determine.

A variety of litter quality indices based on the initial chemical composition have been used as decomposition predictors, such as N concentration or the C/N ratio (Taylor *et al.*, 1989; Zhang and Zak, 1995; Montañez, 1998), the lignin concentration (Meentemeyer, 1978), the lignin-to-N ratio (Melillo *et al.*, 1982), lignin + cellulose, the holocellulose to lignocellulose quotient (McClagherty and Berg, 1987) and P or C/P ratios (Vitousek *et al.*, 1994). Most of these relations have been established in temperate forest, and the types of litter considered do not encompass a large range of biochemical and physical attributes. Therefore, it is not always possible to generalize these indices to other kinds of vegetation. However, the initial C/N as a general index can provide a broad indication of the decomposition potential (Heal *et al.*, 1997).

We found that *C. lundellii* had both the lowest C/N ratio and highest decay rate, in contrast to *M. zapota* with the highest C/N ratio and the lowest decomposition rate. *M. brownei* showed intermediate behavior. Montañez (1998) found that differences between decay rates of tree species in home gardens could likewise be attributed to C/N ratio. Other studies have made similar findings in a variety of species (Tian *et al.*, 1992, Wieder and Lang, 1982). Zhang and Zak (1995) showed that low C/N and high N measures were correlated with higher microbial activity, and consequently, a higher decomposition rate. Comparing the decomposition rates of the local leaf litter samples and the three species shows that *C. lundellii* has a relatively high rate (Figs 6).

In addition to the low C/N ratio, higher total N and P concentrations were displayed by *C. lundellii* in contrast with *M. zapota*. During decomposition carbon is used as an energy source by decomposers while N is assimilated into cell proteins essential for microbial function, thus a higher N concentration in the leaf promotes decomposition (Singh and Gupta, 1977). The effects of a high initial nitrogen concentration are most marked in earlier stages of decomposition, and then in later stages there is little net change in nitrogen concentration as the organic nitrogen becomes more resistant to decomposition (Singh and Gupta, 1977). We found that N concentration increased slightly during the experiment. Increases in the N concentration of leaf litter have been found in other studies as well (Melillo *et al.*, 1982; Babbar and Ewel, 1989). There are several possible reasons for the accumulation on N, none of which were tested in this investigation. Nitrogen may be added by one or more of several mechanisms: fixation, absorption of atmospheric ammonia, throughfall, dust, insect frass, fungal translocation and/or immobilization (Melillo *et al.*, 1982).

Nitrogen is one of the most common factors limiting litter decomposition as it determines the growth and turnover of microbial biomass mineralizing the organic carbon (Heal *et al.*, 1997). Net release or net immobilization can be predicted from the organic material's C/N ratio or N concentration. If the ratio of C/N is below 20 or the N concentration is above 2.5%, N will be released and the material decomposes rapidly. If C/N is much greater than 20, N is likely to be immobilized until decomposition and respiration lower the C/N ratio (Heal *et al.*, 1997). All tree species in the study had C/N ratios above 20, with a tendency to diminish through time and to become more similar (Fig. 3b).

Annual decay rate constants ( $k$ ) described (see also Olson, 1963) the fast, intermediate and slow decomposition of *C. lundellii*, *M. brownei* and *M. zapota* respectively. Simple linear regression analysis with annual decay rate constant ( $k$ ) as a dependent variable and C/N, C/P ratios and N, P concentrations as independent variables showed, that these indices of litter nutrient quality accounted for 84.21%; 87.39% and 74.86%, 78.29% of variance, respectively. Our results confirm that these parameters are good predictors of decomposition rate in a wide range of forest environments.

In tropical forests,  $k$  values are often greater than  $-1.0$ , indicating that leaf litter turnover occurs in a year or less than a year (Alvarez *et al.*, 1992). The  $k$  values obtained for the three species in this work were greater than  $-1.0$ , meaning that decomposition could occur within a year or less. Montañez (1998) also found high values for home garden tree species ranging between  $-3.51$  to  $-4.77$ . Turnover rates for the leaf fraction of litter are also greater than  $-1.0$  in other dry forests. In Chamela, Martinez-Yrizar (1995) determined the  $k$  value of the leaf fraction to be  $-1.39$ , a value in the range of *M. zapota* (also based on fallen leaves). It is important to note that our study was conducted using tissue collected live for two species, rather than naturally senesced leaves. Fallen leaves are likely to decompose more slowly because of changes in N and P content prior to abscission.

The effect of successional phase on decomposition was not statistically significant in terms mass loss, but there was significant difference in decomposition between both young secondary forest and older forests in separate tests (Tukey,  $p < 0.010$ ). Perhaps this lack of significance is due to low number of experimental units (two

plots per successional phase) and sampling units (litterbags per species). However, it could simply be the variability among species in the relative influence of the successional environment. Annual decay rates increased with forest age within each species, although the exact pattern varied across species. These data suggest that succession stage may in fact play a role in determining the decomposition rates of individual species. If decomposition depends on both the chemical composition of the litter and the environment, this study then suggests the optimal environment for decomposition has been reestablished after 13 years but not yet after 3 years of forest development.

### ***In situ* decomposition of local leaf litter**

The local leaf litter decomposition was slightly faster in early successional phases of secondary vegetation than it was in older secondary forests, indicated by higher estimated decay rates. However, this faster decomposition was not correlated with a lower C/N ratio. It was correlated with higher average P concentration throughout the study and with higher initial N concentration. Perhaps slow decomposition of local leaf litter in later phases of secondary vegetation is due to N immobilization. N content increased with time in all phases, but did so more dramatically in the 13 year old stands (by 60%) and 50 year old stands (by 37%) than in the 3 year old stands (by 23%). Given these trends, the N/P ratio of litter is likely to be lower in early successional forests. This would suggest a less P-limited environment for the microbial community.

Brown and Lugo (1990) have suggested that secondary forests are generally less efficient than mature stands in their return of nutrients to the forest floor, regardless

of age, soil type and disturbance history. Vitousek (1984) pointed out that secondary forest accumulates small amounts of nutrients in the litter compartment, but returns large amounts in litterfall. Thus, nutrient turnover in the litter of young forests is actually high and may result in positive feedbacks on production and growth. Data from this study agree with Vitousek, showing higher nutrient concentrations and more rapid decomposition in early successional species (Figs. 2 and 6). As we found, Toky and Ramakrishnan (1983b) found that the percentage of nutrients in the leaves is higher during the first years of secondary forest fallows (1-5 years old) than in older fallows (10-20 years old). They also found that, relative nutrient stocks in the soil decreased with the fallow age and nutrient stock in vegetation increased, meaning that nutrients are relative getting scarce in the soil with fallow age. Thus, decomposition may decrease with successional development because of the dynamics of forest growth as well as the shift in species.

## **CONCLUSION**

Based on our finding that an early successional species (like *C. lundellii*) had faster decay rates, faster decomposition in early successional forest would be expected. This concurs with the results of our *in situ* leaf litter experiment. When comparing mass loss of individual species along a successional gradient, the results appear somewhat contradictory, as there was a tendency toward more rapid decomposition in older, rather than younger, forests. Taken together, our studies suggest that litter quality explains more variability in decomposition rates than does

environment. In fact, the effects of litter quality and environment appear to work in opposite directions along a successional gradient. Enhanced nutrient content favors more rapid decomposition in young forests, and enhanced microbial environments favor more rapid decomposition in older forests.

## REFERENCES

- Alvarez, S. J. And Becerra, E. R. 1996. Leaf decomposition in a Mexican Tropical Rain Forest. *Biotropica* 28,657-667.
- Alvarez, S. F. J., G. R. Sánchez, I. Sánchez-Gallén and J. A. González-Iturbe. 1992. Métodos para el Estudio de la Productividad Primaria y la Descomposición en Comunidades Terrestres. Cuadernos de Ecología. Facultad de Ciencias. UNAM. 52 pp.
- Anderson, J. M. and J. S. I. Ingram. 1993. Tropical Soil Biology and Fertility: A handbook of methods of analysis. CAB International. 220 pp.
- Anderson, J. M. and Swift, M. J. 1983. Decomposition in Tropical Forest. In: Sutton, S. L., Whitmore, T. C. and A. C. Chadwick (Eds.), Tropical Rain Forest. Blackwell Scientific Publications. pp. 287-309.
- AOAC. 1994. Official Methods of Analysis. Association of Official Analytical Chemists. Washington, D.C. USA.
- Babbar, L. I. and Ewel, J. J. 1989. Descomposición del follaje en diversos ecosistemas sucesionales tropicales. *Biotropica* 21, 20-29.
- Brown, S. and Lugo, A. E. 1990. Tropical secondary forests. *Journal of Tropical Ecology* 6, 1-32.



- Coleman, D. C. and Crossley, D. A. 1996. *Fundamentals of Soil Ecology*. Academic Press. University of Georgia, Athens. 204 pp.
- Cuevas, E. 1995. Biology of the belowground system of tropical dry forest. In: Bullock, S. H., H. A. Mooney and E. Medina (Eds.). *Seasonally Dry Tropical Forests*. Cambridge University Press. 450 pp.
- Díaz, G. J. R. 2000. Uso del suelo y transformación de selvas en un ejido de la reserva de la Biosfera de Calakmul, Campeche, México. Master's Thesis. El Colegio de la Frontera Sur. Chetumal, Quintana Roo. México. 27 pp.
- Ende, Carl N. von. 1993. Repeated-Measures Analysis: Growth and other time-Dependent measures. In: Scheiner, S. M. and Gurevitch, J. (Eds). *Design and Analysis of Ecological Experiments*. Chapman and Hall. New York, USA. 445 pp.
- FAO, 1970. El reconocimiento de los suelos de la Península de Yucatán. Italia. Informe técnico 1, ESR:SF/MEX 6, 51 pp.
- Flores, J.S. and Espejel, I. 1994. Tipos de Vegetación de la Península de Yucatán. *Etnoflora Yucatanense*. Fascículo 3, Universidad Autónoma de Yucatán. 135 pp.
- Flores, J. S., E. Ucán, J. Adrews and L. M. Ortega. 1990. Reporte preliminar de la vegetación de la Reserva de la Biosfera de Calakmul, Campeche. UADY, Mérida, Yucatán. Manuscrito.
- García, E. 1973. Modificaciones al sistema de clasificación climática de Köppen. Instituto de Geografía. UNAM. México, D. F. 246 pp.

- García, G. G. and March, I. J. 1990. Elaboración de la Cartografía Temática Básica y Base Geográfica de Datos para la Zona de Calakmul, Campeche. Informe Final. ECOSFERA. 39 pp.
- Harmon, M. K., D. F. Whighan, J. Sexton and I. Olmsted. 1995. Decomposition and mass of woody detritus in the dry tropical forests of the Northeastern Yucatan Peninsula, Mexico. *Biotropica* 27, 305-316.
- Harris, L. E. 1970. Métodos para el análisis químico y la evaluación biológica de alimentos para animales. Center for Tropical Agriculture Feed Composition Project Livestock. University of Florida, USA.
- Heal, O.W.; Anderson, J.M. and Swift, M.J. 1997. Plant Litter Quality and Decomposition: An Historical Overview. In: Cadish, G. and Giller, K.E. (Eds.). *Driven by Nature. Plant Litter Quality and Decomposition*. CABI Publishing. pp 3-30.
- Herborn, J. L. and Congdon, R. A. 1998. Ecosystem dynamics at disturbed and undisturbed sites in North Queensland wet tropical rain forest. III. Nutrient returns to forest floor through litterfall. *Journal of Tropical Ecology* 14, 217-229.
- Lawrence, D. submitted. Regional and local determinants of seasonal litterfall patterns in the dry tropical forests of the Southern Yucatan. *Ecosystems*.
- Lundell, C. L. 1934. Preliminary sketch of the phytogeography of the Yucatan Peninsula. *Carnegie Institute of Washington* 336, 257-321.
- Martínez-Yrizar, A. 1995. Biomass distribution and primary productivity of tropical dry forests. In: Bullock, S. H., H. A. Mooney and E. Medina (Eds.). *Seasonally Dry Tropical Forests*. Cambridge University Press. pp 326-345.

- Mass, J. M. 1995. Conversion of tropical dry forest to pasture and agriculture. In: Bullock, S. H., H. A. Mooney and E. Medina (Eds.). *Seasonally Dry Tropical Forests*. Cambridge University Press. pp 399-422.
- McClaugherty, C. A. and B. Berg. 1987. Hollocellulose, lignin and nitrogen levels as rate-regulating factors in late stages of forests litter decomposition. *Pedobiologia* 30, 101-112.
- Meentemeyer, V. 1978. Macroclimate and lignin control of litter decomposition rates. *Ecology* 59, 465-472.
- Melillo, J. M.; Aber, J. D. and Muratore, J. F. 1982. Nitrogen and lignin control of hardwood leaf litter decomposition dynamics. *Ecology* 63, 621-623.
- Miranda, F. 1958. Estudios acerca de la vegetación en la Península de Yucatán. In: Beltrán, E. (Ed.) *Los Recursos Naturales del Sureste y su Aprovechamiento*. Publ. Inst. Mex. Nat. Renov. Tomo II: 215-271.
- Montañez, E. P. I. 1998. Producción de hojarasca y aporte de nutrimentos en los huertos familiares de Hocabá y Sahcabá, Yucatán, México. Master's Thesis. Facultad de Medicina Veterinaria y Zootecnia. Universidad Autónoma de Yucatán, México. 110 pp.
- Murphy P.G. and Lugo A.E. 1986. Ecology of tropical dry forest. *Annual Review of Ecology and Systematics* 17, 67-88.
- Olson, J. S. 1963. Energy storage and the balance of producers and decomposer in ecological systems. *Ecology* 44, 322-330.
- Prieto, T. D.; Vázquez, G. Ma. M. and Rodríguez, A. C. 1999. Comunidades de la mesofauna edáfica en una selva baja inundable de la Reserva de la Biósfera de Sian Kaan, Quintana Roo, México. *Rev. Biol. Trop.* 47, 489-492.

- Proctor, J. 1983. Tropical forest litterfall. I. Problems of data comparison. In: Sutton, L. S., T. C. Whitmore and A. C. Chadwick. (Eds.), Tropical Rain Forest: Ecology and Management. Blackwell Scientific Publications. pp 267-273.
- Rzedowski, J. 1981. Vegetación de México. Ed. LIMUSA. México. 432 pp.
- Singh, J. S. and Gupta, R. S. 1977. Plant decomposition and soil respiration in terrestrial ecosystems. The Botanical Review 43, 499-528.
- Steel, R.G.D and Torrie, J. H. 1988. Bioestadística: Principios y procedimientos. Ed. McGraw-Hill. Mexico. 622 pp.
- Sundarapandian, S.M. and Swamy, P.S. 1999. Litter production and leaf litter decomposition of selected tree species in tropical forest at Kodoyar in the Western Ghats, India. Forest Ecology and Management 123, 231-244.
- Swift, M. J. and Anderson, J. M. 1989. Decomposition. In: Lieth, H. and Werger, M. J. A. (Eds.), Tropical Rain Forest Ecosystems. Biogeographical and Ecological Studies. Elsevier. pp 547-569.
- Taylor, B; D. Parkinson and W.J. Parsons. 1989. Nitrogen and lignin contents predictors of litter decay rates: a microcosm test. Ecology 70, 97-104.
- Tian, G.; B. T. Kang and L. Brussaard. 1992. Effects of chemical composition on N, Ca and Mg release during incubation of leaves from selected agroforestry and fallow plant species. Biogeochemistry 16, 103-119.
- Toky, O. P. and Ramakrishnan, P. S. 1983a. Secondary succession following slash and burn agriculture in North-Eastern India. I. Biomass, litterfall and productivity. Journal of Tropical Ecology 71,747-757.

- Toky, O. P. and Ramakrishnan, P. S. 1983b. Secondary Succession Following Slash and Burn Agriculture in North-Eastern India. II. Nutrient cycling. *Journal of Tropical Ecology* 71, 747-757.
- Turner, B.L. II, S. Cortina Villar, D. Foster, J. Geoghegan, E. Keys, P. Klepeis, D. Lawrence, P. Macario Mendoza, S. Manson, Y. Ogneva-Himmelberger, D. Perez Salicrup, R. Roy Chowdhury, B. Savitsky, L. Schneider, B. Schmook, C. Vance. *In press*. Deforestation and Agricultural Change in the Southern Yucatán Peninsular Region. *Forest Ecology and Management*.
- Vitousek, P. M. 1984. Litterfall, nutrient cycling, and nutrient limitation in tropical forests. *Ecology* 65, 285-298.
- Vitousek, P. M.; D. R. Turner; W. J. Parton and R. L. Sanford. 1994. Litter decomposition on the Mauna Loa environment matrix, Hawai'i: patterns, mechanisms, and models. *Ecology* 75, 418-429.
- Whigham, D. F., I. Olmsted, E. C. Cano and M. E. Harmon. 1991. The Impact of hurricane Gilbert on trees, litterfall, and woody debris in a dry tropical forest in the North Eastern Yucatan Peninsula. *Biotropica* 23, 434-441.
- Whigham, D. F., P. Z. Towle, C. E. Cabrera, J. O. Neill and Ley, E. 1990. The effect of annual variation in precipitation on growth and litter production in a tropical dry forest in the Yucatan of Mexico. *Tropical Ecology* 3, 23-34.
- Wieder, R. K. and Lang, G. E. 1982. A Critique of the analytical methods used in examining decomposition data obtained from litter bags. *Ecology* 63, 1636-1642.
- Zhang, Q. and Zak, J. C. 1995. Effects of gap size on litter decomposition and microbial activity in a subtropical forest. *Ecology* 76, 2196-2204.

Table 1. Summary of decomposition analysis of three types of litter in three different phases of secondary forest development. The symbol \* mark significant differences.

Source of variation	df	MS	F	p		
between subject						
Age	2	16.44	3.096	0.1864		
Species	2	705.25	132.77	0.001*		
Error	3	5.31				
Source of variation	df	MS	F	p	G-G	H-F
within subject						
Time	5	821.49	277.53	0.001*	0.000	0.000
Time-Age	10	4.650	1.57	0.128	0.167	0.148
Time-Species	10	21.917	7.40	0.001*	0.000	0.000
Error	15	2.962				

Green-House = 0.640

Huynh-Feldt = 0.798

G-G and H-F are Greenhouse and Huynh-Feldt adjusted probabilities respectively based on the respective epsilons.

Table 2. Summary of statistical analysis of local leaf litter on three different phases of secondary forest. The symbol \* mark significant differences.

Source of variation	df	MS	F	p		
between subject						
Age	2	77.15	11.38	0.002*		
Error	3	6.78				
Source of variation	df	MS	F	p	G-G	H-F
within subject						
Time	5	298.02	75.26	0.001*	0.000	0.000
Time-Age	10	2.36	0.596	0.100	0.272	0.244
Error	15	3.96				

Green-House = 0.392

Huynh-Feldt = 0.663

G-G and H-F are Greenhouse and Huynh-Feldt adjusted probabilities respectively based on the respective epsilons




Figure 1. Ombrothermic diagram recorded during 1999-2000 in the Zoh Laguna weather station, 30 km from El Refugio, Campeche, Mexico.




Figure 2. Decomposition rates constant (k) for tree species on three different successional stages of secondary dry forest ( $R^2 > 0.91$ ).





Figure 3. a) Leaf dry mass loss (mean  $\pm$  SD) and b) C/N ratio for tree species in tropical secondary dry forest (days 0 to 35 represent dry season; days 63 to 190 represent rainy season).






Figure 4. The relationship between: C/N, C/P ratios (a, b), total N, P contents (c, d) and annual decomposition rate (k) constant of three types of leaf litter.





Figure. 5. a) Local leaf litter weight loss (mean  $\pm$  SD) and b) C/N ratio of three phases of secondary forest development.




Figure 6. A comparison approach of decomposition rate of three types of leaf litter and local leaf litter in each successional phase of secondary forest.